

1. Evaluate the following integrals.

a. $\int x^2 \cos^2(x^3) dx$

b. $\int e^{3x} \sin(2x) dx$

c. $\int \frac{2x^2 - 3x - 3}{(x-1)(x^2 - 2x + 5)} dx$

d. $\int \frac{dx}{x^4 \sqrt{x^2 - 9}}$

e. $\int_0^{\frac{1}{4}\pi} 4 \sec^4(\vartheta) \tan(\vartheta) d\vartheta$

f. $\int_1^{16} \frac{dx}{(1 + \sqrt[4]{x})\sqrt{x}}$

g. $\int \frac{\log(2x)}{x \log(x)} dx$

2. Evaluate the following improper integrals.

a. $\int_0^1 \frac{\log(x)}{\sqrt{x}} dx$

b. $\int_{-\infty}^{\infty} \frac{dx}{x^2 + 9}$

3. Evaluate the following limits.

a. $\lim_{x \rightarrow \frac{1}{2}\pi} \frac{1 - \sin(x)}{(2x - \pi)^2}$

b. $\lim_{x \rightarrow 2} (\sin(\pi/x))^{\tan(\pi/x)}$

4. Let \mathcal{R} be the region bounded by the graphs of $y = \arcsin(x)$, $y = 0$ and $x = 1$.

a. Evaluate the area of \mathcal{R} .

b. Write an integral which represents the volume of the solid obtained by revolving \mathcal{R} about the line defined by $y = \frac{1}{2}\pi$.

c. Write an integral which represents the volume of the solid obtained by revolving \mathcal{R} about the line defined by $x = 2$.

5. Solve the differential equation

$$x\sqrt{1-y^2} - \sqrt{1-x^2} \frac{dy}{dx} = 0,$$

given that $y(1) = \frac{1}{2}\sqrt{3}$. Express y as a(n explicit) function of x .

6. Find an equation of the curve which passes through the point $(-2, e)$ that has the property that the slope of the tangent line at any of its points is equal to the product of the x and y coordinates of that point.

7. Determine whether the sequence $\{a_n\}$ converges or diverges. If the sequence converges find its limit; otherwise, explain why it diverges.

a. $a_n = (-1)^n \frac{\sqrt{n+3}}{5-3\sqrt{n}}$

b. $a_n = n^2 \cos(1/n) - n^2$

8. Find the sum of the series

$$\sum_{n=2}^{\infty} \frac{\pi + (-2)^n}{3^n}.$$

9. Determine whether the series converges or diverges. Justify all assertions carefully.

a. $\frac{1}{3} - \cos\left(\frac{1}{3}\right) + \frac{1}{9} - \cos\left(\frac{1}{9}\right) + \frac{1}{27} - \cos\left(\frac{1}{27}\right) + \frac{1}{81} - \cos\left(\frac{1}{81}\right) + \dots + 3^{-n} - \cos(3^{-n}) + \dots$

b. $\sum_{n=1}^{\infty} \frac{\arctan(n)}{\sqrt[4]{n^9 + 8}}$

c. $\sum_{n=1}^{\infty} \log(1 + 1/n)$

10. Determine whether each of the following series is absolutely convergent, conditionally convergent or divergent. Justify all assertions carefully.

a. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{n!}{3^n n^n}$

b. $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n^2 + 2 + n}}$

11. Find the interval of convergence of the power series

$$\sum_{n=2}^{\infty} (-1)^{n+1} \frac{(x-2)^n}{n \log(n)}.$$

12. Find the Taylor series of $\cos(x)$ centred at $\frac{1}{2}\pi$.

13. Suppose that the power series

$$f(x) = \sum_{n=1}^{\infty} c_n (x-2)^n$$

converges if $x = -2$ and diverges if $x = -3$.

a. Does the series converge if $x = 6$, or does it diverge, or could it either converge or diverge? Explain.

b. Does the series $\sum c_n$ converge, or does it diverge, or could it converge or diverge? Explain.

c. Show that the series $\sum n c_n$ converges.

1. a. If $y = x^3$ then $dy = 3x^2 dx$, so that

$$\int x^2 \cos^2(x^3) dx = \frac{1}{3} \int \cos^2(y) dy = \frac{1}{6} \int (1 + \sin(2y)) dy$$

$$= \frac{1}{6} x^3 + \frac{1}{12} \sin(2x^3) + a.$$

b. Repeated partial integration (integrating the exponential factor) and absorption gives

$$\int e^{3x} \sin(2x) dx = \frac{1}{3} e^{3x} \sin(2x) - \frac{2}{9} e^{3x} \cos(2x) - \frac{4}{9} \int e^{3x} \sin(2x) dx$$

$$= \frac{1}{13} e^{3x} (3 \sin(2x) - 2 \cos(2x)) + b.$$

c. Resolving the integrand into partial fractions, and then revising the second term gives

$$\int \frac{2x^2 - 3x - 3}{(x-1)(x^2 - 2x + 5)} dx = \int \left\{ \frac{-1}{x-1} + \frac{3x-2}{x^2 - 2x + 5} \right\} dx$$

$$= \int \left\{ \frac{-1}{x-1} + \frac{3x-3}{x^2 - 2x + 5} + \frac{1}{(x-1)^2 + 4} \right\} dx$$

$$= -\log|x-1| + \frac{3}{2} \log(x^2 - 2x + 5) + \frac{1}{2} \arctan\left(\frac{1}{2}(x-1)\right) + c.$$

d. If $y = x^{-1} \sqrt{x^2 - 9}$ then $y^2 = 1 - 9x^{-2}$, $y dy = 9x^{-3} dx$ and $9x^{-2} = 1 - y^2$, so

$$\int \frac{dx}{x^4 \sqrt{x^2 - 9}} = \frac{1}{81} \int \frac{9}{x^2} \cdot \frac{x}{\sqrt{x^2 - 9}} \cdot \frac{9}{x^3} dx = \int (1 - y^2) \cdot \frac{1}{y} \cdot y dy$$

$$= \frac{1}{81} \left(y - \frac{1}{3} y^3 \right) + d = \frac{(3 - y^2)y}{243} + d = \frac{(2x^2 + 9)\sqrt{x^2 - 9}}{243x^3} + d,$$

since $3 - y^2 = 2 + 9x^{-2} = (2x^2 + 9)/x^2$.

e. If $y = \sec(\vartheta)$ then $dy = \sec(\vartheta)\tan(\vartheta)d\vartheta$, $y = 1$ if $\vartheta = 0$ and $y = \sqrt{2}$ if $\vartheta = \frac{1}{4}\pi$, so that

$$\int_0^{\frac{1}{4}\pi} 4 \sec^4(\vartheta) \tan(\vartheta) d\vartheta = \int_1^{\sqrt{2}} 4y^3 dy = y^4 \Big|_1^{\sqrt{2}} = 3.$$

f. If $y = 1 + \sqrt[4]{x}$ then $x = (y-1)^4$, $dx = 4(y-1)^3 dy$ and $\sqrt{x} = (y-1)^2$. Also, $y = 2$ if $x = 1$ and $y = 3$ if $x = 16$, so

$$\int_1^{16} \frac{dx}{(1 + \sqrt[4]{x})\sqrt{x}} = 4 \int_2^3 \frac{(y-1)^3}{y(y-1)^2} dy = 4 \int_2^3 (1 - 1/y) dy = 4(y - \log(y)) \Big|_2^3$$

$$= 4 \log\left(\frac{2}{3}e\right).$$

g. If $y = \log(x)$ then $dy = x^{-1} dx$ and $\log(2x) = y + \log(2)$, so

$$\int \frac{\log(2x)}{x \log(x)} dx = \int \frac{y + \log(2)}{y} dy = y + \log(2) \log|y| + g$$

$$= \log(x) + \log(2) \log|\log(x)| + g.$$

2. a. Partial integration gives

$$\int_0^1 \frac{\log(x)}{\sqrt{x}} dx = \lim_{\epsilon \rightarrow 0^+} 2\sqrt{x} \log(x) \Big|_{\epsilon}^1 - 2 \int_{\epsilon}^1 \frac{dx}{\sqrt{x}} = -4 \lim_{\epsilon \rightarrow 0^+} \sqrt{x} \Big|_{\epsilon}^1 = -4,$$

where the first limit is zero by elementary properties of the logarithm.

b. Using symmetry and a standard integral,

$$\int_{-\infty}^{\infty} \frac{dx}{x^2 + 9} = 2 \int_0^{\infty} \frac{dx}{x^2 + 9} = \frac{2}{3} \lim_{t \rightarrow \infty} \arctan\left(\frac{1}{3}x\right) \Big|_0^t = \frac{2}{3}\pi.$$

3. a. If $y = \frac{1}{2}\pi - x$ then, since $\cos(y) = 1 - \frac{1}{2}y^2 + \frac{1}{24}y^4 - \mathcal{O}(y^6)$, it follows that

$$\lim_{x \rightarrow \frac{1}{2}\pi} \frac{1 - \sin(x)}{(2x - \pi)^2} = \lim_{y \rightarrow 0} \frac{1 - \cos(y)}{4y^2} = \frac{1}{8}.$$

b. If $y = \sin(\pi/x)$ then

$$\lim_{x \rightarrow 2} (\sin(\pi/x))^{\tan(\pi/x)} = \lim_{y \rightarrow 1} y^{\frac{y}{\sqrt{1-y^2}}} = \lim_{y \rightarrow 1} (1 + y - 1)^{\frac{1}{y-1} \cdot \frac{-y\sqrt{1-y}}{\sqrt{1+y}}}$$

$$= e^0 = 1.$$

4. Observe that $\mathcal{R} = \{(x, y): 0 \leq x \leq 1, 0 \leq y \leq \arcsin(x)\}$, or equivalently, $\mathcal{R} = \{(x, y): 0 \leq y \leq \frac{1}{2}\pi, 0 \leq x \leq \sin(y)\}$.

a. The area of \mathcal{R} is equal to

$$\int_0^1 \arcsin(x) dx = x \arcsin(x) \Big|_0^1 - \int_0^1 \frac{x}{\sqrt{1-x^2}} dx = \frac{1}{2}\pi + \sqrt{1-x^2} \Big|_0^1$$

$$= \frac{1}{2}\pi - 1,$$

by partial integration and inspection. Alternatively, the area of \mathcal{R} is equal to

$$\int_0^{\frac{1}{2}\pi} (1 - \sin(y)) dy = (y + \cos(y)) \Big|_0^{\frac{1}{2}\pi} = \frac{1}{2}\pi - 1.$$

b. The solid obtained by revolving \mathcal{R} about the line defined by $y = \frac{1}{2}\pi$ consists of annuli of outer radius $\frac{1}{2}\pi$ and inner radius $\frac{1}{2}\pi - \arcsin(x)$, for $0 \leq x \leq 1$, so its volume is equal to

$$\pi \int_0^1 \left\{ \left(\frac{1}{2}\pi\right)^2 - \left(\frac{1}{2}\pi - \arcsin(x)\right)^2 \right\} dx.$$

Alternatively, the solid consists of concentric cylindrical shells of radius $\frac{1}{2}\pi - y$ and height $1 - \sin(y)$, for $0 \leq y \leq \frac{1}{2}\pi$, so its volume is equal to

$$2\pi \int_0^{\frac{1}{2}\pi} \left(\frac{1}{2}\pi - y\right)(1 - \sin(y)) dy.$$

c. The solid obtained by revolving \mathcal{R} about the line defined by $x = 2$ consists of concentric cylindrical shells of radius $2 - x$ and height $\arcsin(x)$, for $0 \leq x \leq 1$, so its volume is equal to

$$2\pi \int_0^1 (2 - x) \arcsin(x) dx.$$

Alternatively, the solid consists of annuli of inner radius 1 and outer radius $2 - \sin(y)$, for $0 \leq y \leq \frac{1}{2}\pi$, so its volume is equal to

$$\pi \int_0^{\frac{1}{2}\pi} \left\{ (2 - \sin(y))^2 - 1 \right\} dy.$$

5. Separating the variables and integrating gives

$$\frac{1}{\sqrt{1-y^2}} \frac{dy}{dx} = \frac{x}{\sqrt{1-x^2}}, \quad \text{and hence} \quad \arcsin(y) = a - \sqrt{1-x^2},$$

in which $a = \frac{1}{3}\pi$ since $y(1) = \frac{1}{2}\sqrt{3}$. Therefore, $y = \sin\left(\frac{1}{3}\pi - \sqrt{1-x^2}\right)$.

6. The condition on the slope of the tangent line is equivalent to

$$\frac{dy}{dx} = xy, \quad \text{i.e.,} \quad e^{-\frac{1}{2}x^2} \frac{dy}{dx} - e^{-\frac{1}{2}x^2} xy = 0 \quad \text{or} \quad \frac{d}{dx} \left\{ e^{-\frac{1}{2}x^2} y \right\} = 0,$$

Thus, $e^{-\frac{1}{2}x^2} y$ is constant, and therefore $e^{-\frac{1}{2}x^2} y = e^{-\frac{1}{2}(-2)^2} e = e^{-1}$, using the initial condition. So the curve is defined by $e^{-\frac{1}{2}x^2} y = e^{-1}$, or $y = e^{\frac{1}{2}x^2 - 1}$.

7. a. Since

$$\lim_{n \rightarrow \infty} \frac{\sqrt[n]{n+3}}{5-3\sqrt[n]{n}} = -\frac{1}{3}, \quad \text{it follows that} \quad \left\{ (-1)^n \frac{\sqrt[n]{n+3}}{5-3\sqrt[n]{n}} \right\}$$

(oscillates and) diverges.

b. With $x = 1/n$, that $\cos(x) = 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6 + \mathcal{O}(x^8)$, implies that

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \left\{ n^2 (\cos(1/n) - 1) \right\} = \lim_{x \rightarrow 0} \frac{\cos(x) - 1}{x^2} = \frac{1}{2}.$$

8. By a direct computation,

$$\sum_{n=2}^{\infty} \frac{\pi + (-2)^n}{3^n} = \pi \sum_{n=2}^{\infty} \left(\frac{1}{3}\right)^n + \sum_{n=2}^{\infty} \left(-\frac{2}{3}\right)^n = \frac{\pi}{9} \cdot \frac{1}{1 - \frac{1}{3}} + \frac{4}{9} \cdot \frac{1}{1 + \frac{2}{3}}$$

$$= \frac{1}{6}\pi + \frac{4}{15}.$$

9. a. Since $\lim \cos(3^{-n}) \neq 0$ (and $\lim 3^n = 0$), the series in question diverges by the vanishing condition.

b. If $n \geq 1$ then

$$0 < \frac{\arctan(n)}{\sqrt[4]{n^9 + 8}} < \frac{\pi}{2n^{9/4}},$$

so the series in question converges with the p -series $\sum n^{-9/4}$ ($p = \frac{9}{4} > 1$) by the comparison test.

c. If $x > 1$ then $1 - 1/x < \log(x) < x - 1$, so in particular (taking $x = 1 + 1/n$)

$$\frac{1}{n+1} < \log(1 + 1/n) < 1/n \quad \text{for } n \geq 1.$$

Hence, the series in question diverges with the harmonic series (*i.e.*, the p -series with $p = 1$).

10. a. If $a_n = n!/(3^n n^n)$, then $a_n > 0$ if $n \geq 1$ and

$$\lim \frac{a_{n+1}}{a_n} = \lim \frac{(n+1)! 3^n n^n}{n! 3^{n+1} (n+1)^{n+1}} = \lim \frac{1}{3} \left(1 + \frac{1}{n}\right)^{-n} = \frac{1}{3} e^{-1} < 1,$$

so the ratio test implies that the series $\sum (-1)^n a_n$ is absolutely convergent.

b. Since

$$a_n = \frac{1}{\sqrt{n^2 + 2 + n}} > \frac{1}{\sqrt{4n^2 + n}} = \frac{1}{3n} > 0,$$

for $n \geq 1$, the series $\sum a_n$ diverges with the harmonic series. But if $n \geq 1$ then $\sqrt{(n+1)^2 + 2 + n + 1} > \sqrt{n^2 + 2 + n} > 0$, and $\lim \{\sqrt{n^2 + 2 + n}\} = \infty$, so $0 < a_{n+1} < a_n$ for $n \geq 1$ and $\lim a_n = 0$; Hence, the Leibniz test implies that the series $\sum (-1)^n a_n$ is convergent. Therefore, the series $\sum (-1)^n a_n$ is conditionally convergent.

11. If

$$\alpha_n = (-1)^{n+1} \frac{(x-2)^n}{n \log(n)}$$

and $x \neq 2$ then

$$\lim \left| \frac{\alpha_{n+1}}{\alpha_n} \right| = |x-2| \lim \left\{ \frac{n}{n+1} \cdot \frac{\log(n)}{\log(n+1)} \right\} = |x-2|,$$

so the ratio test implies that the series $\sum \alpha_n$ is convergent if $|x-2| < 1$, *i.e.*, $1 < x < 3$, and that $\sum \alpha_n$ is divergent if $x < 1$ or $x > 3$. If $x = 1$ then $\alpha_n = -a_n$, where $a_n = (n \log n)^{-1}$ and if $x = 3$ then $\alpha_n = (-1)^{n+1} a_n$. Since $\sum a_n$ is a divergent logarithmic p -series ($p = 1$), the series $\sum \alpha_n$ diverges if $x = 1$. On the other hand, $(n+1) \log(n+1) > n \log(n) > 0$ if $n \geq 2$ and $\lim(n \log(n)) = \infty$, so $a_n > a_{n+1} > 0$ if $n \geq 2$ and $\lim a_n = 0$. Thus, the Leibniz test implies that $\sum (-1)^{n+1} a_n$ is convergent; *i.e.*, that $\sum \alpha_n$ converges if $x = 3$. Therefore, the radius of convergence of $\sum \alpha_n$ is 1, and its interval of convergence is $(1, 3]$.

12. Since $\cos(x) = \cos(\frac{1}{2}\pi + x - \frac{1}{2}\pi) = -\sin(x - \frac{1}{2}\pi)$, the Maclaurin expansion of the sine function yields

$$\cos(x) = -\sin(x - \frac{1}{2}\pi) = \sum_{k=0}^{\infty} \frac{(-1)^{k+1}}{(2k+1)!} (x - \frac{1}{2}\pi)^{2k+1}.$$

13. The interval of convergence of $f(x)$ contains the interval $[-2, 6)$, is contained in the interval $(-3, 7]$, and may be equal to either of these intervals.

a. The series $f(6)$ may or may not converge, as 6 does not belong to $[-2, 6)$.

b. The series $\sum c_n$ is $f(1)$, which converges, since 1 does belong to $[-2, 6)$.

c. As $|4-2| < |-2-2|$, and $f(-2)$ converges, $f(4) = \sum_{n=1}^{\infty} c_n 2^n$ is absolutely convergent. But if $n \geq 1$, then $0 < n < 2^n$, and hence $0 \leq |nc_n| \leq |c_n 2^n|$. So the comparison test implies that $\sum |nc_n|$ converges with $f(4)$, and therefore the series $\sum nc_n$ is (absolutely) convergent.